

CONSENSUS STATEMENT

**SINGAPORE INTEGRATED
24-HOUR ACTIVITY GUIDELINES
FOR EARLY CHILDHOOD
(< 7 YEARS)**

January 2022



**ACADEMY OF MEDICINE
SINGAPORE**



**COLLEGE OF PAEDIATRICS AND
CHILD HEALTH, SINGAPORE**

INTRODUCTION

Early childhood is a critical period for growth and development, forming the foundation for future and lifelong well-being.(1) Adopting healthy lifestyle behaviours in early childhood can potentially influence and shape behaviours later in life.(2) Frameworks have been developed, such as from the Harvard Centre on the Developing Child, and are advocated for early childhood health promotion and disease prevention;(3) these form key strategies in reducing future non-communicable diseases (NCDs). The World Health Organization's (WHO's) Global Action Plan for the Prevention and Control of Non-Communicable Diseases (2013-2020), highlighted that exposure to risk factors of NCDs often starts in early life and interventions in early childhood often offer the best protection against these NCDs.(4)

The current Singapore guidelines on physical activity for children less than 7 years were updated in 2013.(5) Over recent years, there is emergent evidence surrounding physical activity, sedentary behaviour and sleep for this group of children, and how these concepts relate to one another, within a 24-hour period, for better health outcomes.(6-10) This workgroup also integrated dietary choices and eating behaviours, which are closely linked to movement behaviours in terms of, but is not limited to optimising energy balance, which is important for obesity prevention.(11) Encapsulating all these elements, we developed the Singapore Integrated 24-hour Activity Guidelines for Early Childhood (0 – <7 years).

HEALTH OUTCOMES OF LIFESTYLE BEHAVIOURS

We present the global evidence for health outcomes of physical activity, sedentary behaviour, sleep, and diet and eating habits in children aged 0 to 5 years.

Physical activity provides many health benefits in early childhood, including motor and cognitive development, cardiometabolic, musculoskeletal and psychosocial health.(12) Children in this age group should be encouraged to regularly participate in a variety of activities regularly; those who engage in more physical activity overall and/or at a higher intensity (i.e. moderate to vigorous or vigorous intensity), consistently enjoyed favourable health benefits.(13-15) Tummy time (i.e. prone position) appears to be positively associated with motor development for infants,(16, 17) while outdoor play (under adult supervision) is reported to confer health benefits on preschoolers.(18)

Excessive sedentary behaviour and screen time can have detrimental health effects to children in their early years.(19) Prolonged sedentary screen time is adversely associated with adiposity, poor motor and cognitive development and impaired psychosocial health.(20-22) Indeed, prolonged sitting, reclining or lying are also unfavourably associated with adiposity or motor development.(23, 24)

Sleep is essential for growth and good health in the early years.(25) As a newborn grows, he/she regulates his/her sleep through the establishment of a circadian cycle with less daytime naps and more night-time sleep.(26, 27) Children of different ages require different

optimal sleep durations.(28-30) However, shorter sleep in children could be the result of increased screen time around bedtime and collectively may be associated with higher levels of adiposity, poor growth and emotional dysregulation.(31)

A healthy diet provides optimal nutrition for a child's physical and cognitive development.(32) Nutritional needs and eating patterns change with each progressive stage of the childhood.(33) Setting good eating habits and shaping positive eating behaviours in the early years help form the foundation for a healthy diet, which can reduce the future risks of overweight or obesity, as well as protect against NCDs.(34)

LOCAL STUDIES

The Growing Up in Singapore Towards healthy Outcomes (GUSTO) study is a longitudinal cohort investigation that commenced in 2009, to investigate what influence early development has on body composition and metabolic health.(35) Studies of screen time and sedentary behaviour in the GUSTO cohort showed that the average screen-time for infants (12 months) and toddlers (2 years) was 2.0 and 2.4 hours/day respectively. (36, 37) Screen-time in infants was negatively associated with later cognition (composite IQ and verbal IQ), while for toddlers, higher screen time was associated with less physical activity and greater sedentary behaviour. (36, 38)

The sleep duration of children less than 2 years of age in the GUSTO cohort was significantly associated with body length; shorter sleep duration was also associated with higher body mass index and shorter body length for those at 3 months of age. (39) Infants in the GUSTO cohort who were fed breast milk showed better gross motor skills at 2 years and better cognitive performance at both 2 and 4.5 years of age, when compared to formula-fed infants. (40) Higher intake of sugar sweetened beverages in young children (18 months to 5 years) were associated with higher levels of adiposity and greater risk of overweight or obesity. (41)

A separate cross-sectional study of 78 Singaporean preschoolers, using wrist-worn accelerometers, showed that the children spent a median of 7.8 hours/day in sedentary behaviour and 0.5 hours/day in moderate- to vigorous-intensity physical activity. The same study also revealed that the preschool teachers were not familiar with physical activity guidelines and that parents reported very little outdoor playtime with children after preschool.(42)

GUSTO also reviewed the adherence to (Canada/Australia) 24-hour movement guidelines in 864 children at 5.5 years old and results showed that few children (5.5%) met all of the movement guidelines.(43)

A more recent study examined the proportion of preschoolers meeting the WHO guidelines on physical activity, sedentary behaviour and sleep, and the effect on their quality of life. More than 2000 parents of preschoolers were surveyed and only 9.6% met all of the

recommendations, while 12.6% did not meet any of the recommendations. This study also showed that the health-related quality of life increased as the preschoolers achieved more recommendations.(44) The nationally representative study in terms of ethnicity is apparently the largest prevalence study in Singapore on sleep, physical activity and sedentary behaviour among preschoolers.

AIM OF CONSENSUS STATEMENT

This guidance provides a holistic approach for developing and maintaining good health amongst children in early childhood (i.e. pre-primary schooling age) in Singapore, by integrating physical activity, sedentary behaviour, sleep, and dietary and eating habit advice. It is equally important to understand that these activities are closely related in influencing health outcomes and time-use behaviour, and to organise them within a daily 24-hour period. Incorporating healthy dietary and eating habits with movement behaviours encourage children to practise and adopt these recommended habits and behaviours at a young age thereby conferring good health.

These recommendations are for all healthy infants (0 – <1 year), toddlers (1 – <3 years) and preschoolers (3 – <7 years), regardless of gender, cultural background or socioeconomic status. Children with special needs or medical conditions should consult a qualified medical professional for additional guidance.

METHODS

The consensus workgroup included physicians (neonatologist, paediatrician, sports physician and family physician), allied health professionals (dietitian, exercise physiologist, physiotherapist), academics, educators and researchers from multiple institutions and organisations.

The workgroup assessed the evidence reviews conducted for the WHO Guidelines on Physical Activity, Sedentary Behaviour and Sleep for children under 5 years of age, and the 24-Hour Movement Guidelines for children less than 5 years of age from Canada, Australia and South Africa.(6-9) Relevant evidence for children aged 5 to less than 7 years from WHO Guidelines on Physical Activity and Sedentary Behaviour and the 24-Hour Movement Guidelines for Children and Youth/Young People from Canada and Australia were also reviewed.(45-47) The literature was updated to September 2021 through an electronic search of Medline databases and the keywords used included “infant”, “toddler”, “preschool”, “physical activity”, “sedentary behaviour”, “sleep”, “eating habit” and “diet”. The update included systematic reviews, randomised control trials and cohort studies. Only results in English language were considered. The health outcomes included cardiometabolic health, physical fitness, bone and skeletal health, adiposity, motor and cognitive development, behaviour development and psychosocial health.

The workgroup used the GRADE-ADOLOPMENT approach,(48) which builds on the GRADE Evidence to Decision (EtD) framework,(49) to provide a structured and transparent methodology for healthcare recommendations. It evaluates the strength of recommendations from associated guidelines and the quality of evidence supporting the recommendations. Regular meetings were conducted for workgroup members to discuss and achieve a consensus on the adoption and/or adaption of the WHO guidelines for the local paediatric population.

These recommendations are intended for healthcare professionals providing holistic care to infants, toddlers and preschoolers including education and the promotion of healthy activities that form the foundation for life-long well-being. The full EtD frame work is included as supplementary material.

CONSENSUS STATEMENTS FOR INFANTS (0 – <1 YEAR)

***Physical Activity:* Be physically active several times a day, where more is better, in a variety of forms and within a safe and supervised environment. Activities should include non-screen-based interactive floor-based play and tummy time. For those not yet mobile, tummy time should start soon after birth, building up towards at least 30 minutes spread throughout the day. Planning a daily routine of physical activities may be helpful.**

Physical activity in infants is associated with improved measures of adiposity, motor skill development, psychosocial, and cardiometabolic health indicators.(12) For infants not yet mobile, tummy time, defined as awake prone positioning on a firm surface, is positively associated with multiple developmental aspects.(6, 50) Tummy time has positive effects on global development,(51) particularly gross motor development,(52, 53) body mass index and prevention of brachycephaly.(54, 55) Infants can start on tummy time soon after birth and build up from a few minutes towards at least 30 minutes spread throughout the day. Infants above 3 months of age can aim to achieve a minimum of 1 hour of prone activities spread throughout the day. During tummy time, the infant can be encouraged to play and should be supervised by a responsible adult caregiver.

***Sedentary Behaviour:* Avoid restraining and leaving infants unattended for more than 1 hour at a time. Any form of screen time, including background screen time, is not recommended. When the infant is seated, reclined or lying down, caregivers are encouraged to engage the infant in singing, reading, storytelling and imaginative play. Having a daily routine for activities, sleep and meals may be useful in reducing the amount of sedentary behaviour.**

Infants should not be restrained (e.g. strollers and high chairs) and left unattended for more than 1 hour at a time. Screen time in infants is associated with unfavourable measures of adiposity and decreased scores on measures of psychosocial health and cognitive

development.(19) Screen time in infants confers no benefits and is unfavourably associated with cognitive development,(19) sleep duration and quality,(56, 57) and gross motor development.(58) Any form of screen time, including background screen time which may cause distractions, is not recommended for infants.(6) When the infant is sedentary, engaging in 'serve and return' activities such as reading, singing, storytelling or imaginative play with a caregiver is encouraged.

Sleep: Have a daily total amount of 14-17 hours (for 0-3 months of age) and 12-15 hours (for 4-11 months of age) of sleep, including naps, to promote optimal health. It is recommended for infants to sleep on their back in their own cot, in the same room as their caregivers to ensure sleep safety. Develop a regular sleep time routine to help infants fall asleep with ease.

Infants spend most of their time sleeping, which can be up to 80% in newborns.(59) Good sleep is well known to improve cognitive,(60, 61) physical,(62) and social outcomes,(63) reduce obesity as well as reduce the risk of sudden infant death syndrome.(64-66) Good quality sleep improves family well-being and is an important predictor of maternal health.(67, 68) Although there are cultural differences in sleep duration and practices,(69) you can start developing regular bedtime routines when infants are 2-3 months old and provide a conducive sleep environment to improve sleep duration.(26, 70-74) Good sleep safety practices include infants sleeping supine in their own cot and in the same room as the caregivers.(75-77)

Diet and Eating Habit: Breastfeeding is recommended for infants when possible. From 4 to 6 months of age, introduce a variety of development- and culture-appropriate solid foods of various textures and flavours, that is prepared with no added salt and sugar. Provide a daily routine of having meals spaced 2-3 hours apart in the daytime to avoid overfeeding.

Breast milk supports an infant's nutritional requirements during the first 6 months of life, and provides antibodies to support an infant's health, growth and development.(78, 79) It is recommended that infants are exclusively breastfed for at least the first 6 months of life, with continued use of prescribed medication or vitamin and mineral supplementation if recommended by the physician. Mothers should adhere to food safety and hygiene recommendations if breast milk is expressed and stored. Should human milk be unavailable, infants should be provided with formula milk. In support of healthy bone development, vitamin D supplementation of 400IU per day is recommended for fully and partially breastfed infants due to the low bioavailability in breast milk.(79) With increasing energy and nutrient requirements beyond what breast milk can provide, infants should be started on complementary foods between 4-6 months of age, depending on their developmental readiness.

Iron-containing foods should be encouraged as a first line of defence to prevent deficiency in infants. These can include iron-fortified cereals, pureed meat and poultry, plain tofu or legumes, with textures suited to the infant's stage of development. Salt should not be added to foods for infants as their kidneys are immature and unable to excrete excess salt, thus presenting a safety concern. Food and drinks containing added sugars should be avoided for infants, reducing the risk of dental caries and preventing a learned preference for sugar. Overconsumption of sugar-laden food has been associated with an increased risk of becoming overweight or obese.(41, 79-81) There is no evidence that delaying the introduction of potentially allergenic food prevents food allergies.(79) Therefore, potential allergenic foods such as dairy products, egg, wheat, crustacean shellfish, fish, soy, tree nuts and peanuts should be introduced as part of complementary foods. For infants with a family history of atopy or mild eczema, it is recommended that egg and peanut be introduced one at a time, between 4-6 months of age, once the infant is able to tolerate solid food.(82)

Repeated exposure to a variety of food across all the main food groups (grains and alternatives, lean proteins and alternatives, fruits, and vegetables) is necessary to promote food acceptance and provide infants with the range of required nutrients. Caregivers should strive to recognise an infant's hunger and satiety cues that will support responsive consumption by timely initiation and termination of the feeding process.(82) Evidence reveal non-responsive caregiver feeding practices, such as the use of extremely controlling, restrictive, rewarding or pressure feeding, to be associated with a higher risk of childhood obesity. Thus it is recommended that guidance on responsive feeding be provided to caregivers, so as to promote appropriate weight gain among infants.(83, 84)

CONSENSUS STATEMENTS FOR TODDLERS (1 – <3 YEARS)

Physical Activity: Accumulate at least 180 minutes in a variety of physical activities, where more is better, at any intensity spread throughout the day within a safe environment. Daily outdoor play for toddlers is highly encouraged. Caregivers should actively participate in all forms of physical play with toddlers.

Toddlers should engage a spread of physical activities of light, moderate and vigorous intensities. (12) This should include a variety of physical activities that are fun and encourage exploration, involving movement skills such as walking, running, crawling, climbing, balancing, bending, dancing and playing with balls. The more active play the toddlers achieve, the better. Toddlers who engaged in at least an hour of moving freely each day had significantly stronger object and locomotor skills.(85) Caregivers should participate actively with toddler during both indoor and outdoor play, as such positive interactions are associated with better developmental skills, reduced risk for obesity, and accumulate physical activity.(86, 87)

Structured and unstructured play are important for a toddler's global development and these activities can take place in indoor and outdoor environments.(88) In childcare centres, more

than half of toddler's indoor moderate to vigorous physical activities occur in modifiable open spaces and during class transitions. Enhancing childcare structure quality and inclusion of modifiable open spaces can promote physical activity and reduce sedentary time for toddlers.(89, 90) Being outdoors also increases physical activity, with more playtime and time spent engaged in moderate to vigorous physical activities. In turn, these are associated with better sleep outcomes in toddlers.(57) Furthermore, spending at least 2 hours in outdoor play daily can help prevent the early onset of myopia.

Sedentary Behaviour: Avoid restraining toddlers on a seat for more than 1 hour at a time. Screen time, regardless of the type of device, is not recommended for toddlers younger than 18 months of age, and should be limited to less than 1 hour per day for toddlers 18 months and above. When sitting or lying down, it would be most beneficial to engage the toddler in singing, reading, storytelling or imaginative play.

Sedentary behaviour amongst toddlers, include the use of any screen device, reading, drawing, eating, travelling in a vehicle, whilst sitting, reclining or lying.(91) Prolonged periods of being restrained in a seat or in a supine position is associated with high levels of adiposity and less favourable motor development.(19, 92) Screen-based sedentary behaviours also have unfavourable effects on motor and cognitive development, psychosocial health, social skills (ability to develop positive relationships and interact with others effectively), physical activity and poor sleep outcomes across early childhood.(19, 57) International guidelines consistently recommend that toddlers should not be restrained on their seats for more than 1 hour at a time, and those of less than 18 months of age should have no exposure to screens.(6-8, 93) Even when sedentary, engaging in interactive activities with toddlers such as reading, singing and storytelling has greater potential for cognitive and social development, compared to screen time and solitary activities.(19, 94)

Sleep: Have a daily total amount of 11-14 hours of sleep with regular sleep and wake-up times. Develop a bedtime routine and keep to a consistent bedtime. Provide a conducive sleep environment and avoid screen time before night-time sleep.

Develop a bedtime routine and keep to a consistent bedtime.(95, 96) Provide a conducive sleep environment that is dark, quiet and of comfortable temperature, and avoid screen time 30 minutes before night-time sleep.(97) Recent literature supports that short sleep duration during toddlerhood is associated with greater risk of depressive symptoms and poorer temperament in later childhood.(98, 99) Short sleep duration is also linked to obesogenic eating behaviours.(100-103) Toddlers with short sleep durations also tend to have higher blood pressure later in life. (104)

Regular bedtimes and routines help toddlers to sleep longer and better,(95, 96) and may be important for obesity prevention.(105) Adaptive bedtime activities like storytelling or

cuddling help toddlers sleep longer and have fewer sleep problems.(106) Screens emit blue light that suppresses endogenous melatonin production, in turn resulting in shorter sleep duration, later bedtimes and longer time to fall asleep.(107, 108) Poor quality sleep environment (e.g. crowded, noisy and uncomfortable) is associated with shorter sleep durations, later bedtimes and longer sleep latency.(109, 110)

Diet and Eating Habit: Continue to increase the variety of foods offered to your toddler and wean off milk as the main source of nutrition. Introduce healthy family meals and offer whole milk and water, while establishing a structured routine for meal and snack times. Avoid screen time during meal times. Using food to soothe your toddler or as a reward is discouraged.

Toddlers are reliant on caregivers to establish their feeding habits. These include what, when and how food is consumed.(79, 80, 111, 112) Fresh, minimally-processed foods should be prepared with little or no added sugar and salt, with continual exposure and/or provision of foods across all major food groups that are in unison with healthy family eating habits. As established in the latest Dietary Guidelines for Americans,(79) there is no clear evidence that formula milk should be continued beyond 12 months of age. Pasteurised full cream milk, or fortified unsweetened soy milk, can be incorporated in the toddler's diet from around 12 months of age to meet protein, calcium and vitamin D requirements, accompanied by adequate solid foods.(79, 80) Sugar-sweetened beverages (e.g. juice drinks, sports drinks and regular soft drinks) and caffeinated beverages (e.g. tea, coffee and cola drinks) should not be given before two years of age, and avoided as much as possible thereafter.(79, 113) Instead, plain water should be offered to meet hydration requirements.

Establishing a structured routine for meal and snack times for toddlers is an important component of effective responsive feeding practices, where caregivers also recognise and react to hunger and fullness cues of the toddler.(111, 112) Synthesising findings from randomised-controlled trials, Perez-Escamilla et al. (2017) concluded that responsive feeding practices demonstrated improvements in weight outcomes among toddlers 1 to 2 years of age.(112) Picky eating is also a natural occurrence in the feeding process and toddlers should not be pressured to consume new foods.(79, 80) Instead, they should be provided with regular and frequent exposure to non-preferred foods, increasing familiarity and promoting acceptance. Other toddler behaviours that affect feeding habits such as active play, screen time, sleep and techniques to soothe the toddler, are also influenced by caregivers. Poor sleep routines in the first two years of life, as well as the use of food to soothe the toddler have been associated with poor dietary quality and increased risk of obesity in early childhood,(80) and are therefore not encouraged.

CONSENSUS STATEMENTS FOR PRE-SCHOOLERS (3 – <7 YEARS)

***Physical Activity:* Accumulate at least 180 minutes of physical activity at any intensity spread throughout the day and within a safe environment. At least 60 minutes should be of moderate– to vigorous-intensity, where more is better, and the physical activities can be accumulative and take different forms. Older preschoolers (5 – 6 years of age) should be exposed to a variety of age-appropriate vigorous-intensity play and engage in muscle- and bone-strengthening activities several times a week. Daily outdoor active play among preschoolers is highly encouraged. Caregivers should participate actively with pre-schoolers during all forms of active play.**

Physical activity engagement amongst preschoolers is associated with multiple health benefits, especially when it is moderate- to vigorous-intensity.(12, 114, 115) Evidence supports a positive association between physical activity and motor and cognitive development.(114) Studies have shown that physical activity is associated with favourable adiposity, motor development, physical fitness, psychosocial, cardiometabolic and bone health in preschoolers.(7, 12) In addition, a strong foundation in childhood movement competence is associated with lifelong participation in physical activity; therefore, preschoolers should be encouraged to participate in a variety of activities encompassing fundamental movement skills and age-appropriate/modified sports in a safe environment.(116-118) Evidence from a local study showed that lower primary children failed to demonstrate age-appropriate movement proficiency, indicating a critical need for physical activity interventions at the preschool age.(119)

The high local prevalence of myopia is a serious health concern and daily outdoor play for at least 2 hours can prevent myopia onset.(120, 121) Moreover, outdoor play confers many other learning opportunities for preschoolers, caregivers and educators.(122) Outdoor play with preschoolers provides respite from excessive ‘near work’ (e.g. reading and screen time), and also promotes parent-child bonding and creating meaningful memories for both parents and preschoolers that are enduring.(123)

As preschoolers spend several hours in school on weekdays, there is scope to capitalise more on preschoolers’ natural tendency for movement and physical play. Policies/guidelines should be developed to encourage physical activity throughout the day, such as incorporating it as part of all subject domains, allowing movement breaks hourly and curating learning environments that facilitate preschoolers to move and stay active in childcare centres and kindergartens.(124)

***Sedentary Behaviour:* Limit the total daily amount of sedentary behaviour, such as sitting, reclining or lying down, and take breaks during extended periods of time spent being sedentary. Recreational sedentary screen time, regardless of the type of screen device, should be limited to less than 1 hour per day.**

Recreational screen-based sedentary behaviour such as television viewing and handheld device use is of particular importance.(91) The WHO evidence-based guidelines acknowledged that sedentary behaviour, and in particular recreational screen time among children aged 3-6 years, bore detrimental effects on their fitness, adiposity and behaviour or sleep.(6, 125) The detrimental effects of early-life screen viewing, regardless of the type of screen device, on movement behaviours and adiposity later in life have also been observed in prospective cohort studies among young Singaporean children.(38, 126, 127)

While sedentary behaviour, for instance during educational periods, cannot be completely eliminated, regular movement breaks, such as in the form of active play are essential to minimise adverse health effects.(128) Instead of using screen devices during recreational periods, engaging in reading, drawing, storytelling or imaginative play with a caregiver is encouraged.(19, 94) When engaging in recreational screen time, age-appropriate and engaging content for preschoolers is recommended.

***Sleep:* Have a total of 10 - 13 hours (for 3 - 5 years of age) or 9 - 11 hours (for 6 years of age) daily sleep. Older preschoolers may not need to nap if sufficient sleep has been obtained at night. Develop a bedtime routine and keep to a consistent bed and wake-up time. Provide a conducive sleep environment and avoid screen time before bed.**

Achieving the number of recommended hours of sleep is associated with better health outcomes in terms of physical, psychological, and cognitive well-being. Shorter sleep duration is associated with higher adiposity levels,(129-133) poorer emotional regulation,(134, 135) more screen time,(136-138) higher risk of injuries,(139, 140) poorer cognitive development,(141-144) increased hyperactivity-inattention,(145) reduced physical activity,(137, 146) and poorer quality of life.(147) The total sleep duration, includes both naps and nocturnal sleep, but older preschoolers may not need the former if sufficient has been obtained at night. A bedtime routine should be developed that involves a wind-down period and avoiding screen time 30 minutes before sleep. Aim to maintain this consistency across weekdays and weekends. Providing a conducive sleep environment that is dark, quiet and of comfortable temperature can help preschoolers sleep better.(97)

***Diet and Eating Habit:* Encourage healthy eating habits as a family, with caregivers as role models. Limit the amount and frequency of sugar-sweetened beverage consumption. Provide a structured routine for meal and snack times in appropriate portions that support growth and development. Avoid screen time during meal times. Teach your preschooler to recognise hunger and satiety cues.**

Dietary habits are shaped at a young age and persist later into life. Through continuous positive caregiver modelling, a regular household eating routine provides opportunities for coordinated family meals and regulation of appetite, therefore influencing the overall diet quality of young children.(79, 148, 149) Limiting consumption of sugar-sweetened foods and beverages (including those naturally present in honey, syrups, fruit juices and fruit juice concentrates) to no more than 10% of total energy intake can curb the risk of overweight or obesity and dental caries in children.(41, 79, 81) Similarly, beverages added with non-sugar substitutes such as sugar alcohol (e.g. sorbitol and xylitol) and intense sweeteners (e.g. acesulfame K, sucralose and aspartame) are discouraged as these sweeteners may also cultivate a preference for sweet food and drinks in their diet. Consuming a nutritious breakfast as part of their daily routine is strongly encouraged, as it has been associated with better diet quality and healthy body weight.(149) Structure-based or limit-setting strategies, such as serving appropriate portions, disallowing screen time during family meals, and exerting some caregiver control to moderate preschoolers' intake, can help children develop self-regulation and autonomy in eating behaviours.(79, 148)

Preschoolers are more likely to overeat when watching television or using a screen device during mealtimes and may learn unhealthy food habits from advertisements and programmes. A balance of allowance and control is needed, as being indulgent to a preschooler's food requests may override his or her ability to eat according to internal hunger and satiety cue. Equally, excessive restraint of a preschooler's food intake may unintentionally teach him or her to use food to manage negative emotions.(79, 148) Both of these may lead to unhealthy effects such as overeating and excess weight gain.(79, 148)

CONSENSUS STATEMENTS FOR ALL GROUPS (0 – <7 YEARS)

***Integration:* Aim to achieve most or all recommendations on physical activity, sedentary behaviour, sleep and diet for the best results**

The recommendations for physical activity, sedentary behaviour, sleep and eating habits are closely related in terms of health benefits and making up the 24 hours of a child's day. The greatest health benefits can be achieved by meeting all the recommendations; more physical activity, less sedentary time, longer sleep duration, healthy eating habits and positive dietary choices.(10, 44, 150-152) Equivalent beneficial health outcomes can be obtained by achieving various combinations of the recommendations. Both combinations of

more physical activity with longer sleep duration, and the combination of less sedentary time with longer sleep duration can improve cognitive development and reduce risks of adiposity.(6, 10) Replacing sedentary time with physical activity is associated favourably with fitness and motor development.(6, 10) Healthy and sensible dietary habits promote growth, development and maintenance of a healthy weight.(32, 34)

Conclusion

Recent evidence has shifted the international trend towards integrating physical activity, sedentary behaviour and sleep within a 24-hour period for better health outcomes in young children (i.e. infants, toddlers and preschoolers). The inclusion of diet and eating habits complement these recommendations in supporting growth and development, as well as obesity prevention. Establishing these healthy behaviours in early childhood offers them the best protection against future NCDs. However, local studies have demonstrated that a significant proportion of young children in Singapore do not adopt these recommendations and showed poorer health outcomes. Therefore, it is timely to introduce and promote these guidelines to young children and their caregivers to give them the best start in their lives.

As these young children may be cared for by various caregivers (e.g. parents, grandparents and teachers) and transit through different environments as they grow (e.g. home, infant-care, nursery and kindergarten), there is a role for these recommendations to be adopted as daily habits in family units and also as policies in child-care and preschool centres. In conclusion, infants, toddlers, preschoolers and their caregivers are recommended to adopt all domains of these guidelines to achieve the best health outcomes. Families and schools can start by identifying a domain that they can feasibly embed into their everyday life and aim to achieve all the recommendations in good time.

ACKNOWLEDGEMENTS

This document was developed by Singapore Integrated 24-Hour Activity Guidelines for Early Childhood Study Workgroup which comprised key members from the Singapore community including members from the College of Paediatrics & Child Health of Academy of Medicine Singapore; Singapore Integrated Platform for Research in Advancing Metabolic Health Outcomes in Women and Children (IPRAMHO), led by KK Women's and Children's Hospital (KKH), in partnership with the SingHealth Polyclinics (SHP) and National Healthcare Group Polyclinics (NHGP), Perinatal Society of Singapore and Exercise is Medicine Singapore. The initiative is supported by the research group of IPRAMHO, an NMRC funded joint collaborative pot centre grant of KKH, SHP and NHGP. This multidisciplinary group is initiated by A/Prof Ng Kee Chong and Prof Tan Kok Hian and chaired by Dr Benny Loo.

This guideline summary, produced by the College of Paediatrics & Child Health of Academy of Medicine Singapore, supported by Health Promotion Board (HPB), Singapore & Sport Singapore (SportSG) and partnered with Exercise is Medicine Singapore and Sports Medicine Association Singapore, Perinatal Society of Singapore, Singapore Paediatric Society (SPS), The College of Family Physicians Singapore (CFPS), Singapore Medical Association (SMA), Association for Early Childhood Educators, Singapore (AECES) & Association of Early Childhood and Training Services (ASSETS) acts as an educational aid and reference for healthcare professionals practicing in Singapore. The guideline summary does not define a standard of care, nor is it intended to dictate an exclusive course of management. It presents recognised clinical methods and techniques for consideration by practitioners for incorporation into their practice. It is acknowledged that management may vary and must always be responsive to the need of individual patients, resources, and limitations unique to the institution or type of practice.

WORKGROUP MEMBERS

| Authors: | |
|--|--|
| (1) Dr Benny Kai Guo LOO | <i>Chair, Singapore Integrated 24-Hour Activity Guidelines for Early Childhood Study Workgroup Consultant General Paediatrics Service & Sports Medicine Service KK Women's and Children's Hospital</i> |
| (2) Dr Dinesh SIRISENA | <i>Consultant Sports & Exercise Medicine Khoo Teck Puat Hospital</i> |
| (3) Associate Professor Falk MÜLLER-RIEMENSCHNEIDER | <i>Saw Swee Hock School of Public Health National University of Singapore</i> |
| (4) Professor Michael Yong Hwa CHIA | <i>Physical Education & Sports Science National Institute of Education Nanyang Technological University</i> |

| | |
|--|---|
| (5) Adjunct Associate Professor Benedict Chi'-Loong TAN | <i>Chief & Senior Consultant Department of Sport and Exercise Medicine Changi General Hospital</i> |
| (6) Associate Professor Ngiap Chuan TAN | <i>Family Physician and Senior Consultant SingHealth Polyclinics</i> |
| (7) Clinical Associate Professor Oon Hoe TEOH | <i>Head & Senior Consultant Respiratory Medicine Service KK Women's and Children's Hospital</i> |
| (8) Ethel Jie Kai LIM | <i>Dietician Nutrition & Dietetics KK Women's and Children's Hospital</i> |
| (9) Dr Mohammad Ashik ZAINUDDIN | <i>Head & Senior Consultant Sports and Exercise Medicine Service KK Women's and Children's Hospital</i> |
| (10) Joanne Shumin GAO | <i>Exercise Specialist Sport and Exercise Medicine Service KK Women's and Children's Hospital</i> |
| (11) Dr Poh Chong CHAN | <i>Head & Senior Consultant Division of General Ambulatory Paediatrics and Adolescent Medicine, Department of Paediatrics, Khoo Teck Puat - National University Children's Medical Institute National University Hospital</i> |
| (12) Dr Teresa Shu Zhen TAN | <i>Consultant Division of General Ambulatory Paediatrics and Adolescent Medicine, Department of Paediatrics, Khoo Teck Puat - National University Children's Medical Institute National University Hospital</i> |
| (13) Dr Nirmal KAVALLOOR VISRUTHAN | <i>Consultant Department of Neonatology KK Women's and Children's Hospital</i> |
| (14) Professor Victor Samuel RAJADURAI | <i>Senior Consultant Department of Neonatology KK Women's and Children's Hospital</i> |
| (15) Dr Moira Suyin CHIA | <i>Consultant General Paediatrics Service KK Women's and Children's Hospital</i> |
| (16) Dr Nur Adila Binte Ahmad HATIB | <i>Consultant General Paediatrics Service KK Women's and Children's Hospital</i> |
| (17) Dr Shirong CAI | <i>Principal Investigator Singapore Institute for Clinical Sciences Agency for Science, Technology and Research (A*STAR), Singapore</i> |
| (18) Dr Ju Lynn ONG | <i>Senior Research Fellow Centre for Sleep and Cognition Yong Loo Lin School of Medicine National University of Singapore</i> |

| | |
|---|--|
| (19) Associate Professor June Chi Yan LO | <i>Assistant Professor Centre for Sleep and Cognition Yong Loo Lin School of Medicine National University of Singapore</i> |
| (20) Associate Professor Mary Foong Fong CHONG | <i>Assistant Professor Saw Swee Hock School of Public Health National University of Singapore</i> |
| (21) Dr Le Ye LEE | <i>Senior Consultant Department of Neonatology, Khoo Teck Puat - National University Children's Medical Institute National University Hospital</i> |
| (22) Dr Elaine Chu Shan CHEW | <i>Senior Consultant Adolescent Medicine Service KK Women's and Children's Hospital</i> |
| (23) Dr Ratnaporn SIRIAMORNSARP | <i>Family Physician Tampines Polyclinic SingHealth Polyclinics</i> |
| (24) Dr Miriam LEE | <i>Senior Manager Division of Children & Youth Programme Development Sport Singapore</i> |
| (25) Dr Aaron SIM | <i>Assistant Director Division of Physical Activity & Weight Management Health Promotion Board, Singapore</i> |
| (26) Jean ANG | <i>IPRAMHO Assistant Manager Division of Obstetrics & Gynaecology KK Women's and Children's Hospital</i> |
| (27) Dr Elaine Phaik Ling QUAH | <i>IPRAMHO Senior Research Fellow Division of Obstetrics & Gynaecology KK Women's and Children's Hospital</i> |
| (28) Clinical Associate Professor Kee Chong NG | <i>Senior Consultant Children's Emergency KK Women's and Children's Hospital</i> |
| (29) Professor Kok Hian TAN | <i>Head & Senior Consultant Perinatal Audit & Epidemiology Unit Department of Maternal Fetal Medicine KK Women's and Children's Hospital</i> |
| (30) Dr Chui Mae WONG | <i>Senior Consultant Department of Child Development and Neonatology KK Women's and Children's Hospital</i> |
| (31) Shelly-Anne SHERWOOD | <i>Senior Physiotherapist KK Women's and Children's Hospital</i> |
| (32) Dr Siao Hui TOH | <i>Senior Physiotherapist KK Women's and Children's Hospital</i> |
| (33) Professor Yung Seng LEE | <i>Senior Consultant, Division of Paediatric Endocrinology and Diabetes, Khoo Teck Puat-National University Children's Medical Institute, National University Hospital</i> |

First version published 21 January 2022

College of Paediatrics & Child Health of Academy of Medicine Singapore

SINGAPOREAN INTEGRATED 24-HOUR ACTIVITY GUIDELINES FOR EARLY CHILDHOOD

CONSENSUS STATEMENTS FOR INFANTS (0 – < 1 YEAR)

Physical Activity: Be physically active several times a day, where more is better, in a variety of forms and within a safe and supervised environment. Activities should include non-screen-based interactive floor-based play and tummy time. For those not yet mobile, tummy time should start soon after birth, building up towards at least 30 minutes spread throughout the day. Planning a daily routine of physical activities may be helpful.

Sedentary Behaviour: Avoid restraining and leaving infants unattended for more than 1 hour at a time. Any form of screen time, including background screen time, is not recommended. When the infant is seated, reclined or lying down, caregivers are encouraged to engage the infant in singing, reading, storytelling and imaginative play. Having a daily routine for activities, sleep and meals may be useful in reducing the amount of sedentary behaviour.

Sleep: Have a total amount of 14 - 17 hours (for 0 - 3 months of age) and 12 - 15 hours (for 4 - 11 months of age) of daily sleep, including naps, to promote optimal health. It is recommended for infants to sleep on their back in their own cot, in the same room as their caregivers to maintain sleep safety. Develop a regular sleep time routine to help infants fall asleep easier.

Diet and Eating Habit: Breastfeeding is recommended for infants when possible. From 4 to 6 months of age, introduce a variety of development- and culture-appropriate solid foods of various textures and flavours, that is prepared with no added salt and sugar. Provide a daily routine of having meals spaced 2-3 hours apart in the daytime to avoid overfeeding.

CONSENSUS STATEMENTS FOR TODDLERS (1 – < 3 YEARS)

Physical Activity: Accumulate at least 180 minutes in a variety of physical activities, where more is better, at any intensity spread throughout the day within a safe environment. Daily outdoor play for toddlers is highly encouraged. Caregivers should actively participate in all forms of physical play with toddlers.

Sedentary Behaviour: Avoid restraining toddlers on a seat for more than 1 hour at a time. Screen time, regardless of the type of device, is not recommended for toddlers younger than 18 months of age, and should be limited to less than 1 hour per day for

toddlers 18 months and above. When sitting or lying down, it would be most beneficial to engage the toddler in singing, reading, storytelling or imaginative play.

Sleep: Have a daily total amount of 11-14 hours of sleep with regular sleep and wake-up times. Develop a bedtime routine and keep to a consistent bedtime. Provide a conducive sleep environment and avoid screen time before night-time sleep.

Eating Habit and Diet: Continue to increase the variety of foods offered to your child and wean off milk as the main source of nutrition. Introduce healthy family meals and offer whole milk and water, while establishing a structured routine for meal and snack times. Using food to soothe your child or as a reward is discouraged.

CONSENSUS STATEMENTS FOR PRE-SCHOOLERS (3 – < 7 YEARS)

Physical Activity: Accumulate at least 180 minutes of physical activity at any intensity spread throughout the day and within a safe environment. At least 60 minutes should be of moderate- to vigorous-intensity, where more is better, and the physical activities can be accumulative and take different forms. Older preschoolers (5 – 6 years of age) should be exposed to a variety of age-appropriate vigorous-intensity play and engage in muscle- and bone-strengthening activities several times a week. Daily outdoor active play among preschoolers is highly encouraged. Caregivers should participate actively with preschoolers during all forms of active play.

Sedentary Behaviour: Limit the total daily amount of sedentary behaviour, such as sitting, reclining or lying down, and take breaks during extended periods of time spent being sedentary. Recreational sedentary screen time, regardless of the type of screen device, should be limited to less than 1 hour per day.

Sleep: Have a daily total of 10-13 hours (for 3-5 years of age) or 9-11 hours (for 6 years of age) sleep. Older preschoolers may not need to nap if sufficient sleep has been obtained at night. Develop a bedtime routine and keep to a consistent bed and wake-up time. Provide a conducive sleep environment and avoid screen time before bed.

Diet and Eating Habit: Encourage healthy eating habits as a family, with caregivers as role models. Limit the amount and frequency of sugar-sweetened beverage consumption. Provide a structured routine for meal and snack times in appropriate portions that support growth and development. Avoid screen time during meal times. Teach your preschooler to recognise hunger and satiety cues.

CONSENSUS STATEMENTS FOR ALL GROUPS (0 – < 7 YEARS)

Integration: Aim to achieve most or all recommendations on physical activity, sedentary behaviour, sleep and diet for the best results.

PRACTICAL REFERENCE FOR ACTIVITIES FOR EARLY CHILDHOOD

Physical Activity

Physical activity for early childhood refers to any activity that gets children moving. All activity counts! This guide provides examples of activities that children in the early years can participate in to accumulate physical activity time throughout the day.

For infants (0 – <1 year), physical activity includes tummy time and floor-based play that are appropriate for their developmental milestones. For toddlers (1 – <3 years) and pre-schoolers (3 – <7 years), physical activity should include a **wide variety of movement experiences, in different environments**. This involves activities both **indoors and outdoors**, which take place in various settings **in school and out of school**. **Daily outdoor active play** is highly encouraged as it provides opportunities for toddlers and preschoolers to develop **fundamental movement skills** and to explore their environment.

Types of Physical Activities for Infants

For infants, physical activity includes being carried in an upright position, reaching up to swipe and swat toys when lying in supine, elevating legs off the surface to kick, play in side-lying, rolling from back to tummy and reverse, tummy time, reaching across the body to play when in sitting, crawling, pulling to stand and cruising along furniture. These activities are encouraged on a firm surface and gradually progressed to dynamic surfaces as the infant becomes more confident with the skill. Many of these activities can be incorporated into day-to-day caregiving activities.

It is especially important to remember to limit the number of hours that infants spend in containers or being restraint e.g. rockers, strollers that inhibit their freedom of movement. When infants are allowed to move, they are able to discover their bodies, explore their environment and this further motivates them to learn and move.

| Activities | Tips for parents/caregivers |
|--|--|
| Carried in an upright position (this works on infant's head and trunk control) | <ul style="list-style-type: none">• Try carrying infant over your shoulder when burping them after milk time.• At two months, carry infant facing out, you can allow infant to rest the back of their head against your chest if still wobbly, when more stable, slide infant's buttocks up your chest so that their head is not touching your chest. If you use a carrier, face them out when awake. |
| Reaching up to swipe and swat | <ul style="list-style-type: none">• After changing infant's nappy, when lying on their back, allow infant the opportunity to reach up to swipe and swat at toys. Start with 1-2 minutes and gradually progress to 5-10 minutes.• If infant is unable to reach upwards, used a folded blanket under infant's shoulders and upper arm, one on each side. |

| | |
|--|---|
| Elevating legs off the surface to kick | <ul style="list-style-type: none"> • After changing infant's nappy, dangle a toy at infant's feet and encourage upward kicking of legs. • You may fold a blanket and place under infant's buttocks. This will further help to engage infant's core muscles during kicking. |
| Play in side-lying (to engage the side abdominal muscles - obliques) | <ul style="list-style-type: none"> • When infant is awake, perhaps whilst getting their milk ready or doing your household chores – place infant on a firm surface, lying on their side. You may use a bolster or a rolled bath towel behind their back to prevent them from rolling back. Use a book, contrast cards, or a small toy in their hand to engage them. Alternate sides. |
| Rolling | <ul style="list-style-type: none"> • When changing infant's nappy, instead of lifting legs up, bend infant's hips and knees up to 90 degrees and roll infant to the side to remove the soiled nappy. Then back again when placing the new one. |
| Reaching in sitting | <ul style="list-style-type: none"> • Once infant can sit independently for a few seconds, try placing their favourite toy in front of them so that they reach for it. Progress this by then placing the toy at the sides, encouraging them to reach across their body to get the toy. Consider a cause-and-effect toy or a container where they can put things into and take them out. |
| Crawling | <ul style="list-style-type: none"> • Crawling is initially best practiced on a firm play mat. Ensure the environment is safe and large enough to promote exploration. • Progress crawling to varied surfaces and then over pillows and cushions. |
| Pulling to stand | <ul style="list-style-type: none"> • Start this activity where infant can grip onto something to pull self-up (e.g. cot rails). • Progress this to places where gripping would be harder (e.g. sofa or a wall). |
| Cruising along the furniture | <ul style="list-style-type: none"> • Start this activity where infant can grip onto something, use a toy to encourage side stepping along the furniture. • Progress by making small gaps between the furniture. |

Tips on Tummy Time for Infants

- Your infant may start tummy time as soon as they are brought home.
- Place your infant on their tummy when they are awake, relaxed and rested. Place them on a firm surface like a mat or on your chest so that they may see your face. Start with 1-2 minutes per session and lengthen to 5-10 minutes a few times a day as your infant grows and becomes stronger. Face your infant whilst talking or singing to them. If they appear tired or distressed, roll them onto their backs for a rest and try again later.
- If your infant finds it consistently difficult to lift their head when in tummy time, use your hand to place on their buttocks, this can give them more stability and allow them to push up a little longer.
- When your infant is strong enough, they may try to roll over from their back to their tummy. You may place one or two toys on a firm surface around your infant to encourage this.
- You may place the infant's favourite toy just out of reach to encourage them to reach for or creep towards the toy.

- Make tummy time part of your infant’s daily routine, for example for a short duration before or after diaper changes.

Types of Physical Activities for Toddlers and Preschoolers

Toddlers and preschoolers can be engaged in a wide variety of physical activities through a range of **fundamental movement skills (FMS)**. FMS underpin daily living activities and provide the foundation for participation in sports and other forms of complex movement skills as the toddler/preschooler grow up. The early years are critical for establishing this foundation.

Parents/caregivers play an important role in enhancing the FMS of their toddler/preschooler at home and in school via facilitation, motivation and personal participation. Parents/Caregivers can also facilitate learning of FMS by providing simple cues, demonstration and playing with their toddler/preschooler.

FMS are generally categorised into three main themes:

1. **Locomotor Skills** refers to body movement from one location to another. Many locomotor skills are used daily (e.g. running after a bus, leaping over a puddle), in many games and sports (e.g. jumping up to catch a ball) and during active play (e.g. crawling through a tunnel or climbing in the playground). Other examples of locomotor skills include walking, sliding, hopping, and skipping.
2. **Object Control (manipulative) Skills** require the toddler/preschooler to control an object using part of the body or using an equipment. Object control skills involve:
 - Propulsive skills – Sending an object away (e.g. throwing or kicking a ball)
 - Receptive skills – Receiving an object (e.g. catching or dribbling a ball)
3. **Stability (non-locomotor) Skills** involve a toddler/preschooler in maintaining and/or attaining balance. Stability is a key element for every human movement and necessary for all locomotor and object skills. Examples of stability skills include static and dynamic balance, bending and curling, turning, twisting, and stretching.

(Source: Fun Start Move Smart!: Fundamental movement skills for growing active learners)

Other types of physical activities to encourage participation of preschoolers include swimming, cycling and other forms of modified sport or age-appropriate multi-sport.

Older preschoolers should progress towards participating in a range of physical activities that involve age-appropriate energetic play. They are recommended to carry out moderate to vigorous physical activity (MVPA) for 60 minutes, which is a third of the recommended total daily physical activities (PAs) of 180 minutes.

MVPA should include frequent periods of energetic and dynamic play, spread throughout the day. Moderate-intensity activity refers to activity that make the child breathe harder and heart beats faster than at rest or sitting, such as walking to and from school. For vigorous-intensity activity, a useful guide is to observe if the child is huffing and puffing during the activity, or is not able to say more than a few words without pausing to catch a breath, examples include, playing tag (running or chasing playmates), ball games, skating or rapid cycling. Across an exertion scale of 10, these activities will score about 7 or 8. Examples of

MVPA for the preschooler will include, playing tag (running or chasing playmates), ball games, skating or rapid cycling.

In addition to MVPA, preschoolers should also incorporate muscle and bone strengthening activities. These activities require the preschoolers to bear and lift their own body weight to work against a resistance. Examples include running, jumping, climbing activities such as scaling playground obstacles, skipping rope, dancing and playing games such as hopscotch. Dedicated resistance or weight training regimes are not necessary.

Physical activities for the toddler and preschooler may be organised broadly into two categories: i) structured physical activity with planned objectives and focus, and ii) unstructured physical activity that allows for “free” and unguided play. Both genres are synergistic (i.e. deliberate vs. self-regulated way of achieving activity goals) and should be incorporated into the toddler’s/preschooler’s daily physical activity routine. For example, a Physical Education session with structured and unstructured play covers lesson for a preschooler to balance on a beam and freedom to play alone or with others at a playground.

| Physical Activity Out of School | | |
|--|---|--|
| Different Environments to engage in Physical Activity | Tips for Parents/Caregivers | Suggested Activities/Examples |
| At Home (Indoors) | Create a home environment that encourages movement and exploration | <ul style="list-style-type: none"> • Set up safe spaces for movement (e.g. an area for physical play) • Position furniture in a way that allows child to move freely and encourages movement around the house • House rules that encourage safe movement (e.g. a toddler is allowed to climb up and down low furniture or sofa) • Devote spaces in the house for active play (e.g. corridor for shuttle run, hopscotch, underarm rolling to a target) • Make accessible a bucket of foam balls of different sizes for children to play with • Paste pictures of favourite cartoon characters on the wall for child to aim and throw at • Hang balloon or streamers from the doorway to encourage child to jump, reach and strike • Set up obstacle course that incorporates various activities such as balancing, jumping, throwing and etc. for child to play and navigate around |

| | | |
|---|---|---|
| | Make use of household/recycled material for activities | <ul style="list-style-type: none"> • Create an obstacle course or a fort with cardboard boxes, pillows, chairs, and bedsheets to crawl through or jump over • Rolled up socks, crushed newspaper or soft toys for throw and catch games • Make DIY equipment (e.g. paper plate and short stick as an implement to send and receive a balloon to and fro) • Masking tape on the floor to balance on the line (dynamic balance) |
| | Play with your child (parent-child games) | <ul style="list-style-type: none"> • Hide and seek and treasure hunt (use different locomotor skills) • Keep the balloon in the air with different body parts • Pillow fights • Simon Says • Animal movements • Alphabet poses • A game of Twister • Musical statue • Dance to music |
| | Involve child in household chores (even if a helper is available) | <ul style="list-style-type: none"> • Clean up after meals (e.g., bring plate back to kitchen or wipe the dining table) • Tidy the house (e.g., put toys/books back, return items after use) • Fold clothes and put it back in wardrobe • Water plants |
| On-the-go (Outdoors/Indoors) | Integrate as part of daily activity and commute | <ul style="list-style-type: none"> • Walk up and down stairs • Walk, cycle or scoot to and from school |
| Playground/Play gardens (Outdoors) | Make it a routine for child to visit the playground; Encourage unstructured play | <p>Playgrounds present numerous opportunities for children to explore and develop FMS. Some examples include:</p> <ul style="list-style-type: none"> • Balancing on a balance beam, a log or on an uneven surface • Climbing up a rock wall or rope ladders • Crawling through a tunnel • Jumping/hopping into coloured markers on the floor or on the trampoline • Sliding also involves stability skills such as bending, and transfer of weight to get down the slide |
| | Facilitate active play: Introduce games resulting in moderate to vigorous physical activity (MVPA) | <p>Introduce games with simple rules and play with your child:</p> <ul style="list-style-type: none"> • Create an obstacle course to challenge your child • Catching or Tag, Freeze-Tag, The Floor is Lava (use different locomotor skills) |

| | | |
|---|---|--|
| Outdoor Spaces (e.g., basketball court, void deck, open spaces, grass fields) | Identify available outdoor open spaces in your neighbourhood for your child to participate in MVPA | <ul style="list-style-type: none"> • Games with simple rules: Catching or Tag, Freeze-Tag, What’s the Time Mr Wolf? (use different locomotor skills) • Ball games (e.g. throw and catch, kicking, bat and ball, badminton) • Bubbles (run, jump and strike or catch) • Scooter, Cycle (tricycle, balance bike, two-wheel bike), Roller skate |
| Parks/ Nature Parks/Beach (Outdoors) | Immerse your child in a new environment to engage in active play over the weekend | <ul style="list-style-type: none"> • Nature walks or hike • Walk, run, jump, hop, skip bare feet on different surfaces (e.g., grass, sand) • Walk on uneven ground • Balance on logs • Build sandcastles • Water play/ Wade in the water |
| Sport Facilities (e.g., stadiums, swimming pools, badminton courts, climbing gym) (Outdoors/Indoors) | Let your child experience a variety of sports according to his/her interest | <ul style="list-style-type: none"> • Participating in a range of sports with an age-appropriate programme helps develop FMS. Some examples include: <ul style="list-style-type: none"> ○ Football ○ Athletics ○ Gymnastics ○ Basketball ○ Tennis ○ Hockey ○ Martial arts ○ Swimming ○ Rock climbing |

Physical Activity in School

As toddlers and preschoolers spend many hours in school on weekdays (especially those in childcare centres), preschool educators play a significant role in getting toddlers/preschoolers to be active in physical activity throughout the day. Besides classes that directly involve movement (e.g. Music and Movement, Motor Skills Development), educators should provide other opportunities for toddlers/preschoolers to increase body movements throughout the day. Some suggestions include:

- **Modifying the physical environment** to increase movement of the toddlers/preschoolers as part of their daily routine. Introduce
 - **Active Navigation Routes** – Design spaces and pathways linking different areas of the school using floor markers; install attractive signage to encourage toddler/preschooler to jump, hop, balance etc. to get from one part of the school to another.
 - **Active Play Corners** (e.g., Giant game board that gets toddler/preschooler to perform various FMS)

- Incorporate **movement as part of core curriculum** (e.g., numeracy with movement)
- Introduce **outdoor learning** or **MVPA in a different environment**
- Introduce **Brain breaks** after every hour of sedentary time or when toddlers/preschoolers are restless. This involves quick and simple activities to get toddlers/preschoolers out of their seats to move (e.g., stretching, action songs, animal movements)

Physical Activities within a Preschoolers' Day

There are many ways in which a preschooler is capable of fulfilling the physical activity recommendations on any given day. The following suggestions and examples aim to illustrate a typical school day and a weekend.

On weekdays:

- **Preschool:** Over 2 hours, 1 hour can be structured physical education class, and the remaining 1 hour covers other curriculum such as music or dance class, active involvement in story-telling
- **Home (1 hour)**
 - Playground after school (as part of daily routine, 45-60min)
 - Walk to school, climb up/down stairs (20min)
 - Active play at home.

On weekends

- Beach/Park (picnic, walking in nature, cycling); park connector
- Organised sport with modified games
- Playing at activity facilities (playground, water-park, beach)

Resources

Tummy time

1. <https://polyclinic.singhealth.com.sg/Documents/GrossMotorSkills.pdf>

Play activities for infants

1. <https://polyclinic.singhealth.com.sg/Documents/3month%20DA.pdf>
2. <https://polyclinic.singhealth.com.sg/Documents/6month%20DA.pdf>
3. https://polyclinic.singhealth.com.sg/Documents/12month%20DA_v2.pdf
4. <https://www.healthhub.sg/live-healthy/2030/activity-ideas-to-get-tot-moving>

Outdoor play for toddlers & preschoolers

1. <https://activeparents.myactivesg.com/activities/ap/lets-go-play-outside-5-unique-playgrounds-around-singapore-to-train-your-childs-fundamental-movement-skill> [Video]
2. <https://www.ecda.gov.sg/Educators/Pages/Outdoor-Learning.aspx>

Active Play for preschoolers

1. <https://activeparents.myactivesg.com/activities/ap/active-family-series-school-holiday-activity-pack-2020>
2. <https://activeparents.myactivesg.com/activities/ap/word-up-challenges-with-my-1st-coach> [Video]

3. <https://activeparents.myactivesg.com/activities/ap/father-child-exercises> [Video]
4. <https://activeparents.myactivesg.com/activities/ap/mother-child-exercises> [Video]
5. <https://activeparents.myactivesg.com/parent-child-exercises> [Video]

ActiveSG programmes for preschoolers

1. <https://www.myactivesg.com/programmes/academy>
2. <https://www.activeparents.myactivesg.com>

Active screen-based physical activity for preschoolers

1. <https://raisingchildren.net.au/preschoolers/play-learning/screen-time-healthy-screen-use/screen-time-physical-activity>

Sedentary Behaviour

Parents, caregivers, and teachers can help young children to reduce their sedentary behaviour and to engage in active lifestyles. Consider the following:

| | |
|----------------|---|
| Infant | <ul style="list-style-type: none"> • Talk and sing gently to your infant whilst holding them. When talking, try using a changing tone of voice and vary your expressions. • Read books that have simple, repetitive words and clear pictures to your infant, especially those with rhymes or songs. • Stimulate your infant with toys that are colourful or make noise, encouraging them to reach out for the toy. Encourage your infant to explore toys of different textures (soft, hard, rough) for exploratory play. Household objects like cups, spoons or boxes can be used. Blocks are useful for open-ended play. • Play peek-a-boo with your infant with your hands or another object like a book or napkin. • You may bring your infant outdoors to explore their surroundings. • When playing with your infant, follow their lead and change activities based on their needs. • Avoid restraining your baby in containers e.g. strollers and rockers for prolonged periods. |
| Toddler | <ul style="list-style-type: none"> • Playtime is a powerful way of showing love and connecting with your toddler. • Allow your toddler to take on unstructured, free play. It enables them to develop creativity and learn at their own pace. • A device cannot replace you. Your interaction with your toddler will benefit them extensively. • For toddlers >18 months old, if you allow screen time, consider setting consistent rules and limits (e.g. less than 1 hour of screen time per day). Ensure access to high quality educational programs for your toddlers • Co-view and share screen time with your toddler and talk about what he/she is watching. • Face-to-face interactions is much preferred, which include talking, reading and singing. |

| | |
|--------------------|---|
| | <ul style="list-style-type: none"> • Avoid screen device use while eating meals, in the hour before bedtime, playing with toys or during family interactions. • Parents should be aware of their own screen usage and avoid using screens while spending time with your toddler. |
| Preschooler | <ul style="list-style-type: none"> • Active play and physical activity are important, and curb sedentary time when possible. • When sedentary behaviour cannot be avoided, break up extended periods of sitting or lying down with frequent breaks for movement. • Encourage a variation of postures and avoid sustained positions to avoid muscular aches and pain. • While sedentary, engage in interactive activities together, such as storytelling, playing games and singing. • Minimize the amount of screen viewing as much as possible. • If screen time cannot be avoided, set rules and boundaries and consider setting a family media plan for screen usage, ensure age-appropriate content, avoid screen device use while eating meals, in the hour before bedtime and playing with toys or during family conversations. • Remove TV's and other screen devices from the preschooler's bedroom. • Monitor media content and select high-quality programs appropriate to the preschooler's age, engage in conversations with the preschooler about the media content and co-view with them as much as possible. • Parents should be aware of their own screen usage and avoid using screens while spending time with your toddler. |

Resources

Play activities for infants

1. <https://polyclinic.singhealth.com.sg/Documents/3month%20DA.pdf>
2. <https://polyclinic.singhealth.com.sg/Documents/6month%20DA.pdf>
3. https://polyclinic.singhealth.com.sg/Documents/12month%20DA_v2.pdf
4. <https://www.healthhub.sg/live-healthy/2030/activity-ideas-to-get-tot-moving>

Movement/Fun activities at home

1. <https://www.nuh.com.sg/our-services/Specialties/Paediatrics/Documents/Activity%20Resource%20Book%20-%20PLAY.pdf>
2. <https://www.nuh.com.sg/our-services/Specialties/Paediatrics/Documents/Movement%20activities%20at%20home%20for%20your%20child.pdf>

Advice about screen time

1. <https://www.nuh.com.sg/our-services/Specialties/Paediatrics/Documents/NUH%20CDU%20Screen%20Time%20E-Brochure.pdf>
2. <https://www.healthychildren.org/English/family-life/Media/Pages/Healthy-Digital-Media-Use-Habits-for-Babies-Toddlers-Preschoolers.aspx>
3. <https://childmind.org/article/media-guidelines-for-kids-of-all-ages/>
4. <https://raisingchildren.net.au/preschoolers/play-learning/screen-time-healthy-screen-use/managing-screen-time-3-11-years>

5. <https://raisingchildren.net.au/toddlers/play-learning/screen-time-media/screen-time>

Family media plan

1. <https://www.healthychildren.org/English/media/Pages/default.aspx>

Sleep

Parents, caregivers, and teachers can help young children to meet the sleep recommendations. Consider the following:

| | |
|-----------------------|--|
| <p>Infant</p> | <ul style="list-style-type: none"> • Set a consistent, calming bedtime routine to establish infant’s sleeping behaviours. A regular pre-bedtime plan can include bathing, infant massage, swaddling and dimming lights. • Learn to look out for body language indicating tiredness and need for sleep (e.g. rubbing eyes, arching back and infant not focusing). • Put infant to bed while drowsy but not completely asleep. Avoid nursing infant to sleep after the first few months, to allow infant to learn to fall asleep independently. • If infant is unsettled, provide rhythmic patting to calm infant, but not to the point of sleeping. • If infant remains restless, providing a pacifier may help. However, avoid carrying infant or rocking infant to sleep. • Provide infant opportunities to self soothe when he wakes up. Your infant might fuss or cry before finding a comfortable position and falling asleep. • Allow sufficient daytime naps, as infant may not sleep well at night if overtired. |
| <p>Toddler</p> | <ul style="list-style-type: none"> • Develop a bedtime routine that involves a wind-down period, which helps to relax and anticipate bedtime (e.g. reading a book, a relaxing shower, listening to music). • Avoid screen time (e.g. smart phones, tablets, computers) 30 minutes before sleep. • Avoid naps close to bedtime as it may prevent the toddler from falling asleep or staying asleep. • Set and maintain a consistent bedtime across weekdays and weekends. • Prioritise your toddler’s sleep over other activities – practise good time management to ensure there is enough time for bedtime routine and that the toddler gets the recommended number of hours of sleep. • Provide a conducive sleep environment that is dark, quiet and of comfortable temperature. • Keep the bed only for sleep and rest. • Avoid stimulating and vigorous activities such as exercise and consumption of caffeinated food (e.g. soft drinks, chocolates) or beverages before bedtime. • May consider providing the toddler with an item that the toddler feels secure with (e.g. a favourite pillow, blanket or stuffed toy) . |

| | |
|--------------------|--|
| Preschooler | <ul style="list-style-type: none"> • Develop a bedtime routine that involves a wind-down period. • Avoid screen time 30 minutes before sleep. • Set and maintain a consistent bedtime across weekdays and weekends. • Prioritise your preschooler’s sleep over other activities – practise good time management to ensure there is enough time for bedtime routine and that the preschooler gets the recommended number of hours of sleep. • Provide a conducive sleep environment that is dark, quiet and of comfortable temperature. • Keep the bed only for sleep and rest. • Avoid stimulating and vigorous activities such as exercise and consumption of caffeinated food (e.g. soft drinks, chocolates) or beverages before bedtime. • Be a sleep role model for your preschooler, i.e. sleep sufficiently yourself, during sleep time phones should be on silent mode, switched off, or not be brought into the bedroom. |
|--------------------|--|

| |
|---|
| Resources |
| <p>Helping infant sleep</p> <ol style="list-style-type: none"> 1. https://www.healthhub.sg/live-healthy/1660/helping-baby-sleep 2. https://www.healthxchange.sg/news/bedtime-options-for-babies 3. https://polyclinic.singhealth.com.sg/Documents/GoodSleepHabitsBabies.pdf 4. https://www.healthhub.sg/live-healthy/1195/baby-time-to-unwind 5. https://www.healthhub.sg/live-healthy/1936/how-can-i-get-my-baby-to-sleep-well-and-safely <p>Tips on good sleep</p> <ol style="list-style-type: none"> 1. https://www.healthhub.sg/programmes/117/goodsleep |

Eating Habit and Diet

Parents, caregivers, and teachers can help young children to meet the eating habit and diet recommendations. Consider the following:

| |
|---|
| Nutrition for Breastfeeding Mothers |
| <p>It is important to have a nutritionally-balanced diet, and of adequate portions, as this may affect the quality and quantity of breastmilk. Nutritional requirements during breastfeeding can be found on:</p> <ol style="list-style-type: none"> 1. Integrated Maternal and Child Wellness Hub (SingHealth Polyclinics) web link: <ul style="list-style-type: none"> • https://polyclinic.singhealth.com.sg/Documents/NutritionDuringBreastfeeding.pdf 2. SingHealth YouTube channel web link: <ul style="list-style-type: none"> • https://www.youtube.com/watch?v=NXEDIKOD0Is&list=PLwKZdOHmwfHG_SfEKd_pAjjarcQEvrXKbY&index=7 |

| Nutrition for Young Children | | |
|---|---|---|
| Nutrition-related information for parents | <p>For information on breastfeeding, childhood nutrition and recipes according to various age groups. It includes other information on general care, growth and development, sleep and activity, as well as parenting tips.</p> <p>Web link:</p> <ul style="list-style-type: none"> • https://www.healthhub.sg/programmes/183/parent-hub • https://www.youtube.com/watch?v=Zu-0WnjRzA8&list=PLwKZdOHmwfHG_SfEKdpApjarcQEvrXKbY&index=8 • https://activeparents.myactivesg.com/edu-series/ap/3g-diet-for-preschoolers • https://activeparents.myactivesg.com/edu-series/ap/5-ways-to-get-your-child-to-love-veggies | <p>Search in browser:</p> <p>“Parent Hub HPB”</p> <p>“Singhealth Baby Feeding Tips”</p> <p>“Get Your Kids Going, Growing & Glowing – the 3G Diet for Pre-schoolers”</p> <p>“5 Ways To Get Your Child To Love Veggies”</p> |
| Early Childhood Nutrition | <p>For information on early nutrition according to the following age groups:</p> <ul style="list-style-type: none"> - 0 to 4 months - 4 to 6 months - 6 to 12 months - 12 to 24 months <p>Resource guides include recommended amount for each food group per day and recipes. Contact of helplines and support groups available.</p> | <p>QR code:</p>  |

REFERENCES

1. Shonkoff JP, Boyce WT, McEwen BS. Neuroscience, molecular biology, and the childhood roots of health disparities: building a new framework for health promotion and disease prevention. *JAMA*. 2009;301(21):2252-9.
2. Janz KF, Burns TL, Levy SM, Iowa Bone Development S. Tracking of activity and sedentary behaviors in childhood: the Iowa Bone Development Study. *Am J Prev Med*. 2005;29(3):171-8.
3. Mistry KB, Minkovitz CS, Riley AW, Johnson SB, Grason HA, Dubay LC, et al. A new framework for childhood health promotion: the role of policies and programs in building capacity and foundations of early childhood health. *Am J Public Health*. 2012;102(9):1688- 96.
4. World Health Organization. Global action plan for the prevention and control of noncommunicable diseases 2013-2020. Geneva: World Health Organization; 2013.
5. Health Promotion Board Singapore. National Physical Activity Guidelines - Children and Youth Aged up to 18 years 2013 [Available from: https://www.academia.edu/10443994/National_Physical_Activity_Guidelines_for_Children_and_Youth].
6. World Health Organization. Guidelines on Physical Activity, Sedentary Behaviour and Sleep for Children under 5 Years of Age. Guidelines on Physical Activity, Sedentary Behaviour and Sleep for Children under 5 Years of Age. WHO Guidelines Approved by the Guidelines Review Committee. Geneva2019.
7. Tremblay MS, Chaput JP, Adamo KB, Aubert S, Barnes JD, Choquette L, et al. Canadian 24-Hour Movement Guidelines for the Early Years (0-4 years): An Integration of Physical Activity, Sedentary Behaviour, and Sleep. *BMC Public Health*. 2017;17(Suppl 5):874.
8. Okely AD, Ghersi D, Hesketh KD, Santos R, Loughran SP, Cliff DP, et al. A collaborative approach to adopting/adapting guidelines - The Australian 24-Hour Movement Guidelines for the early years (Birth to 5 years): an integration of physical activity, sedentary behavior, and sleep. *BMC Public Health*. 2017;17(Suppl 5):869.
9. Draper CE, Tomaz SA, Biersteker L, Cook CJ, Couper J, de Milander M, et al. The South African 24-Hour Movement Guidelines for Birth to 5 Years: An Integration of Physical Activity, Sitting Behavior, Screen Time, and Sleep. *J Phys Act Health*. 2020;17(1):109-19.
10. Kuzik N, Poitras VJ, Tremblay MS, Lee EY, Hunter S, Carson V. Systematic review of the relationships between combinations of movement behaviours and health indicators in the early years (0-4 years). *BMC Public Health*. 2017;17(Suppl 5):849.
11. Fernandez-Alvira JM, De Bourdeaudhuij I, Singh AS, Vik FN, Manios Y, Kovacs E, et al. Clustering of energy balance-related behaviors and parental education in European children: the ENERGY-project. *Int J Behav Nutr Phys Act*. 2013;10:5.
12. Carson V, Lee EY, Hewitt L, Jennings C, Hunter S, Kuzik N, et al. Systematic review of the relationships between physical activity and health indicators in the early years (0-4 years). *BMC Public Health*. 2017;17(Suppl 5):854.
13. Butte NF, Puyau MR, Wilson TA, Liu Y, Wong WW, Adolph AL, et al. Role of physical activity and sleep duration in growth and body composition of preschool-aged children.

Obesity (Silver Spring). 2016;24(6):1328-35.

14. Leppanen MH, Nystrom CD, Henriksson P, Pomeroy J, Ruiz JR, Ortega FB, et al. Physical activity intensity, sedentary behavior, body composition and physical fitness in 4-year-old children: results from the ministop trial. *Int J Obes (Lond)*. 2016;40(7):1126-33.
15. Harvey NC, Cole ZA, Crozier SR, Kim M, Ntani G, Goodfellow L, et al. Physical activity, calcium intake and childhood bone mineral: a population-based cross-sectional study. *Osteoporos Int*. 2012;23(1):121-30.
16. De Kegel A, Peersman W, Onderbeke K, Baetens T, Dhooge I, Van Waelvelde H. New reference values must be established for the Alberta Infant Motor Scales for accurate identification of infants at risk for motor developmental delay in Flanders. *Child Care Health Dev*. 2013;39(2):260-7.
17. Dudek-Shriber L, Zelazny S. The effects of prone positioning on the quality and acquisition of developmental milestones in four-month-old infants. *Pediatr Phys Ther*. 2007;19(1):48-55.
18. Brussoni M, Gibbons R, Gray C, Ishikawa T, Sandseter EB, Bienenstock A, et al. What is the Relationship between Risky Outdoor Play and Health in Children? A Systematic Review. *Int J Environ Res Public Health*. 2015;12(6):6423-54.
19. Poitras VJ, Gray CE, Janssen X, Aubert S, Carson V, Faulkner G, et al. Systematic review of the relationships between sedentary behaviour and health indicators in the early years (0-4 years). *BMC Public Health*. 2017;17(Suppl 5):868.
20. Yilmaz G, Demirli Caylan N, Karacan CD. An intervention to preschool children for reducing screen time: a randomized controlled trial. *Child Care Health Dev*. 2015;41(3):443-9.
21. Lin LY, Cherng RJ, Chen YJ, Chen YJ, Yang HM. Effects of television exposure on developmental skills among young children. *Infant Behav Dev*. 2015;38:20-6.
22. Tomopoulos S, Dreyer BP, Berkule S, Fierman AH, Brockmeyer C, Mendelsohn AL. Infant media exposure and toddler development. *Arch Pediatr Adolesc Med*. 2010;164(12):1105-11.
23. Sijtsma A, Sauer PJ, Stolk RP, Corpeleijn E. Infant movement opportunities are related to early growth--GECKO Drenthe cohort. *Early Hum Dev*. 2013;89(7):457-61.
24. Reilly JJ, Armstrong J, Dorosty AR, Emmett PM, Ness A, Rogers I, et al. Early life risk factors for obesity in childhood: cohort study. *BMJ*. 2005;330(7504):1357.
25. Colten HR, Altevogt BM. Sleep Disorders and Sleep Deprivation: An Unmet Public Health Problem. In: Colten HR, Altevogt BM, editors. *Sleep Disorders and Sleep Deprivation: An Unmet Public Health Problem*. The National Academies Collection: Reports funded by National Institutes of Health. Washington (DC)2006.
26. Bathory E, Tomopoulos S. Sleep Regulation, Physiology and Development, Sleep Duration and Patterns, and Sleep Hygiene in Infants, Toddlers, and Preschool-Age Children. *Curr Probl Pediatr Adolesc Health Care*. 2017;47(2):29-42.
27. Iglowstein I, Jenni OG, Molinari L, Largo RH. Sleep duration from infancy to adolescence: reference values and generational trends. *Pediatrics*. 2003;111(2):302-7.

28. Paruthi S, Brooks LJ, D'Ambrosio C, Hall WA, Kotagal S, Lloyd RM, et al. Recommended Amount of Sleep for Pediatric Populations: A Consensus Statement of the American Academy of Sleep Medicine. *J Clin Sleep Med*. 2016;12(6):785-6.
29. Chaput JP, Dutil C, Sampasa-Kanyinga H. Sleeping hours: what is the ideal number and how does age impact this? *Nat Sci Sleep*. 2018;10:421-30.
30. Hirshkowitz M, Whiton K, Albert SM, Alessi C, Bruni O, DonCarlos L, et al. National Sleep Foundation's updated sleep duration recommendations: final report. *Sleep Health*. 2015;1(4):233-43.
31. Chaput JP, Gray CE, Poitras VJ, Carson V, Gruber R, Birken CS, et al. Systematic review of the relationships between sleep duration and health indicators in the early years (0-4 years). *BMC Public Health*. 2017;17(Suppl 5):855.
32. World Health Organization. Healthy Diet Geneva2020 [updated 29 April 2020. Available from: <https://www.who.int/news-room/fact-sheets/detail/healthy-diet>.
33. Health Promotion Board Singapore. Early Childhood Nutrition [Available from: <https://www.healthhub.sg/programmes/122/early-nutrition-for-babies/>].
34. McGuire S. World Health Organization. Comprehensive Implementation Plan on Maternal, Infant, and Young Child Nutrition. Geneva, Switzerland, 2014. *Adv Nutr*. 2015;6(1):134-5.
35. Soh SE, Tint MT, Gluckman PD, Godfrey KM, Rifkin-Graboi A, Chan YH, et al. Cohort profile: Growing Up in Singapore Towards healthy Outcomes (GUSTO) birth cohort study. *Int J Epidemiol*. 2014;43(5):1401-9.
36. Aishworiya R, Cai S, Chen HY, Phua DY, Broekman BFP, Daniel LM, et al. Television viewing and child cognition in a longitudinal birth cohort in Singapore: the role of maternal factors. *BMC Pediatr*. 2019;19(1):286.
37. Bernard JY, Padmapriya N, Chen B, Cai S, Tan KH, Yap F, et al. Predictors of screen viewing time in young Singaporean children: the GUSTO cohort. *Int J Behav Nutr Phys Act*. 2017;14(1):112.
38. Chen B, Bernard JY, Padmapriya N, Ning Y, Cai S, Lanca C, et al. Associations between early-life screen viewing and 24 hour movement behaviours: findings from a longitudinal birth cohort study. *Lancet Child Adolesc Health*. 2020;4(3):201-9.
39. Zhou Y, Aris IM, Tan SS, Cai S, Tint MT, Krishnaswamy G, et al. Sleep duration and growth outcomes across the first two years of life in the GUSTO study. *Sleep Med*. 2015;16(10):1281-6.
40. Pang WW, Tan PT, Cai S, Fok D, Chua MC, Lim SB, et al. Nutrients or nursing? Understanding how breast milk feeding affects child cognition. *Eur J Nutr*. 2020;59(2):609-19.
41. Quah PL, Kleijweg J, Chang YY, Toh JY, Lim HX, Sugianto R, et al. Association of sugar-sweetened beverage intake at 18 months and 5 years of age with adiposity outcomes at 6 years of age: the Singapore GUSTO mother-offspring cohort. *Br J Nutr*. 2019;122(11):1303-12.

42. Chen B, Waters CN, Compier T, Uijtdewilligen L, Petrunoff NA, Lim YW, et al. Understanding physical activity and sedentary behaviour among preschool-aged children in Singapore: a mixed-methods approach. *BMJ Open*. 2020;10(4):e030606.
43. Chen B, Bernard JY, Padmapriya N, Yao J, Goh C, Tan KH, et al. Socio-demographic and maternal predictors of adherence to 24-hour movement guidelines in Singaporean children. *Int J Behav Nutr Phys Act*. 2019;16(1):70.
44. Chia MY, Tay LY, Chua TB. Quality of life and meeting 24-h WHO guidelines among preschool children in Singapore. *Early Childhood Education Journal*. 2020;48(3):313-23.
45. Chaput JP, Willumsen J, Bull F, Chou R, Ekelund U, Firth J, et al. 2020 WHO guidelines on physical activity and sedentary behaviour for children and adolescents aged 5-17 years: summary of the evidence. *Int J Behav Nutr Phys Act*. 2020;17(1):141.
46. Tremblay MS, Carson V, Chaput JP. Introduction to the Canadian 24-Hour Movement Guidelines for Children and Youth: An Integration of Physical Activity, Sedentary Behaviour, and Sleep. *Appl Physiol Nutr Metab*. 2016;41(6 Suppl 3):iii-iv.
47. Okely AD, Ghersi D, Loughran SP, Cliff DP, Shilton T, Jones RA. Australian 24-hour movement guidelines for children (5-12 years) and young people (13-17 years): an integration of physical activity, sedentary behaviour. Canberra, Australian Government. 2019.
48. Schunemann HJ, Wiercioch W, Brozek J, Etzendorf-Ikobaltzeta I, Mustafa RA, Manja V, et al. GRADE Evidence to Decision (EtD) frameworks for adoption, adaptation, and de novo development of trustworthy recommendations: GRADE-ADOLEPMENT. *J Clin Epidemiol*.
49. Moberg J, Oxman AD, Rosenbaum S, Schunemann HJ, Guyatt G, Flottorp S, et al. The GRADE Evidence to Decision (EtD) framework for health system and public health decisions. *Health Res Policy Syst*. 2018;16(1):45.
50. Hewitt L, Kerr E, Stanley RM, Okely AD. Tummy Time and Infant Health Outcomes: A Systematic Review. *Pediatrics*. 2020;145(6).
51. Senju A, Shimono M, Tsuji M, Suga R, Shibata E, Fujino Y, et al. Inability of infants to push up in the prone position and subsequent development. *Pediatr Int*. 2018;60(9):811-9.
52. Kuo YL, Liao HF, Chen PC, Hsieh WS, Hwang AW. The influence of wakeful prone positioning on motor development during the early life. *J Dev Behav Pediatr*. 2008
53. Russell DC, Kriel H, Joubert G, Goosen Y. Prone positioning and motor development in the first 6 weeks of life. *South African Journal of Occupational Therapy*. 2009;39(1):11-4.
54. Koren A, Kahn-D'angelo L, Reece SM, Gore R. Examining Childhood Obesity From Infancy: The Relationship Between Tummy Time, Infant BMI-z, Weight Gain, and Motor Development-An Exploratory Study. *J Pediatr Health Care*. 2019;33(1):80-91.
55. Aarnivala H, Vuollo V, Harila V, Heikkinen T, Pirttiniemi P, Holmstrom L, et al. The course of positional cranial deformation from 3 to 12 months of age and associated risk factors: a follow-up with 3D imaging. *Eur J Pediatr*. 2016;175(12):1893-903.
56. Chen B, van Dam RM, Tan CS, Chua HL, Wong PG, Bernard JY, et al. Screen viewing behavior and sleep duration among children aged 2 and below. *BMC Public Health*.

2019;19(1):59.

57. Janssen X, Martin A, Hughes AR, Hill CM, Kotronoulas G, Hesketh KR. Associations of screen time, sedentary time and physical activity with sleep in under 5s: A systematic review and meta-analysis. *Sleep Med Rev.* 2020;49:101226.
58. Hauck JL, Felzer-Kim IT. Time Spent in Sedentary Activity Is Related to Gross Motor Ability During the Second Year of Life. *Percept Mot Skills.* 2019;126(5):753-63.
59. Georgoulas A, Jones L, Laudiano-Dray MP, Meek J, Fabrizi L, Whitehead K. Sleep-wake regulation in preterm and term infants. *Sleep.* 2021;44(1).
60. Tham EK, Schneider N, Broekman BF. Infant sleep and its relation with cognition and growth: a narrative review. *Nat Sci Sleep.* 2017;9:135-49.
61. Jiang F. Sleep and Early Brain Development. *Ann Nutr Metab.* 2019;75 Suppl 1:44-54.
62. El Halal CDS, Nunes ML. Sleep and weight-height development. *J Pediatr (Rio J).* 2019;95 Suppl 1:2-9.
63. Ben-Zion H, Volkovich E, Meiri G, Tikotzky L. Mother-Infant Sleep and Maternal Emotional Distress in Solo-Mother and Two-Parent Families. *J Pediatr Psychol.* 2020;45(2):181-93
64. Cook F, Conway LJ, Giallo R, Gartland D, Sciberras E, Brown S. Infant sleep and child mental health: a longitudinal investigation. *Arch Dis Child.* 2020;105(7):655-6
65. Franco P, Montemitro E, Scaillet S, Groswasser J, Kato I, Lin JS, et al. Fewer spontaneous arousals in infants with apparent life-threatening event. *Sleep.* 2011;34(6):733-43.
66. McNamara F, Sullivan CE. Sleep-disordered breathing and its effects on sleep in infants. *Sleep.* 1996;19(1):4-12.
67. Marinelli KA, Ball HL, McKenna JJ, Blair PS. An Integrated Analysis of Maternal-Infant Sleep, Breastfeeding, and Sudden Infant Death Syndrome Research Supporting a Balanced Discourse. *J Hum Lact.* 2019;35(3):510-20.
68. Oyetunji A, Chandra P. Postpartum stress and infant outcome: A review of current literature. *Psychiatry Res.* 2020;284:112769.
69. Yu X, Quante M, Rueschman M, Ash T, Kaplan ER, Guo N, et al. Emergence of racial/ethnic and socioeconomic differences in objectively measured sleep-wake patterns in early infancy: results of the Rise & SHINE study. *Sleep.* 2021;44(3).
70. Mindell JA, Telofski LS, Wiegand B, Kurtz ES. A nightly bedtime routine: impact on sleep in young children and maternal mood. *Sleep.* 2009;32(5):599-606.
71. Ragni B, De Stasio S, Barni D, Gentile S, Giampaolo R. Parental Mental Health, Fathers' Involvement and Bedtime Resistance in Infants. *Ital J Pediatr.* 2019;45(1):134.
72. Mindell JA, Williamson AA. Benefits of a bedtime routine in young children: Sleep, development, and beyond. *Sleep Med Rev.* 2018;40:93-108.
73. Paul IM, Hohman EE, Loken E, Savage JS, Anzman-Frasca S, Carper P, et al. Mother-Infant Room-Sharing and Sleep Outcomes in the INSIGHT Study. *Pediatrics.* 2017;140(1).

74. Volkovich E, Bar-Kalifa E, Meiri G, Tikotzky L. Mother-infant sleep patterns and parental functioning of room-sharing and solitary-sleeping families: a longitudinal study from 3 to 18 months. *Sleep*. 2018;41(2)
75. Colson ER, Geller NL, Heeren T, Corwin MJ. Factors Associated With Choice of Infant Sleep Position. *Pediatrics*. 2017;140(3).
76. Goodstein MH, Ostfeld BM. Improvements in Infant Sleep Position: We Can Do Better! *Pediatrics*. 2017;140(3).
77. Task Force On Sudden Infant Death S. SIDS and Other Sleep-Related Infant Deaths: Updated 2016 Recommendations for a Safe Infant Sleeping Environment. *Pediatrics*. 2016;138(5).
78. World Health Organization. Essential nutrition actions: mainstreaming nutrition through the life-course. Geneva: World Health Organization; 2019
79. Phillips JA. Dietary Guidelines for Americans, 2020-2025. *Workplace Health Saf*. 2021;69(8):395.
80. Riley LK, Rupert J, Boucher O. Nutrition in Toddlers. *Am Fam Physician*. 2018;98(4):227-33.
81. World Health Organization. Guideline: Sugars Intake for Adults and Children. Guideline: Sugars Intake for Adults and Children. WHO Guidelines Approved by the Guidelines Review Committee. Geneva 2015.
82. Academy of Medicine Singapore. Consensus statement: prevention of allergy in at-risk infants. Singapore: Academy of Medicine Singapore; 2019 December.
83. Spill MK, Callahan EH, Shapiro MJ, Spahn JM, Wong YP, Benjamin-Neelon SE, et al. Caregiver feeding practices and child weight outcomes: a systematic review. *Am J Clin Nutr*. 2019;109(Suppl_7):990S-1002S.
84. Lindsay AC, Sitthisongkram S, Greaney ML, Wallington SF, Ruengdej P. Non-Responsive Feeding Practices, Unhealthy Eating Behaviors, and Risk of Child Overweight and Obesity in Southeast Asia: A Systematic Review. *Int J Environ Res Public Health*. 2017;14(4).
85. Barnett LM, Hnatiuk JA, Salmon J, Hesketh KD. Modifiable factors which predict children's gross motor competence: a prospective cohort study. *Int J Behav Nutr Phys Act*. 2019;16(1):129.
86. Hager ER, Tilton NA, Wang Y, Kapur NC, Arbaiza R, Merry BC, et al. The home environment and toddler physical activity: an ecological momentary assessment study. *Pediatr Obes*. 2017;12(1):1-9.
87. Jayasuriya A, Williams M, Edwards T, Tandon P. Parents' perceptions of preschool activities: exploring outdoor play. *Early Educ Dev*. 2016;27(7):1004-17.
88. Lee RLT, Lane S, Brown G, Leung C, Kwok SWH, Chan SWC. Systematic review of the impact of unstructured play interventions to improve young children's physical, social, and emotional wellbeing. *Nurs Health Sci*. 2020;22(2):184-96.
89. Zhang Z, Kuzik N, Adamo KB, Ogden N, Goldfield GS, Okely AD, et al. Associations

Between the Child Care Environment and Children's In-Care Physical Activity and Sedentary Time. *Health Educ Behav.* 2021;48(1):42-53.

90. Fees BS, Fischer E, Haar S, Crowe LK. Toddler activity intensity during indoor free-play: stand and watch. *J Nutr Educ Behav.* 2015;47(2):170-5.
91. Tremblay MS, Aubert S, Barnes JD, Saunders TJ, Carson V, Latimer-Cheung AE, et al. Sedentary Behavior Research Network (SBRN) - Terminology Consensus Project process and outcome. *Int J Behav Nutr Phys Act.* 2017;14(1):75.
92. Bruijns BA, Truelove S, Johnson AM, Gilliland J, Tucker P. Infants' and toddlers' physical activity and sedentary time as measured by accelerometry: a systematic review and meta-analysis. *Int J Behav Nutr Phys Act.* 2020;17(1):14.
93. Reid Chassiakos YL, Radesky J, Christakis D, Moreno MA, Cross C, Council On C, et al. Children and Adolescents and Digital Media. *Pediatrics.* 2016;138(5).
94. Carson V, Lee EY, Hesketh KD, Hunter S, Kuzik N, Predy M, et al. Physical activity and sedentary behavior across three time-points and associations with social skills in early childhood. *BMC Public Health.* 2019;19(1):27.
95. Covington LB, Rogers VE, Armstrong B, Storr CL, Black MM. Toddler Bedtime Routines and Associations With Nighttime Sleep Duration and Maternal and Household Factors. *J Clin Sleep Med.* 2019;15(6):865-71.
96. Prokasky A, Fritz M, Molfese VJ, Bates JE. Night-to-Night Variability in the Bedtime Routine Predicts Sleep in Toddlers. *Early Child Res Q.* 2019;49:18-27.
97. Hoban TF. Sleep and its disorders in children. *Semin Neurol.* 2004;24(3):327-40.
98. Sivertsen B, Harvey AG, Reichborn-Kjennerud T, Ystrom E, Hysing M. Sleep problems and depressive symptoms in toddlers and 8-year-old children: A longitudinal study. *J Sleep Res.* 2021;30(1):e13150.
99. Bastien L, Tetreault E, Bernier A. Disentangling the Direction of Associations between Sleep and Temperament in Toddlers. *Behav Sleep Med.* 2020;18(4):523-36.
100. Miller AL, Miller SE, LeBourgeois MK, Sturza J, Rosenblum KL, Lumeng JC. Sleep duration and quality are associated with eating behavior in low-income toddlers. *Appetite.* 2019;135:100-7.
101. Goetz AR, Beebe DW, Peugh JL, Mara CA, Lanphear BP, Braun JM, et al. Longer sleep duration during infancy and toddlerhood predicts weight normalization among high birth weight infants. *Sleep.* 2019;42(2).
102. Zhang Z, Pereira JR, Sousa-Sa E, Okely AD, Feng X, Santos R. The cross-sectional and prospective associations between sleep characteristics and adiposity in toddlers: Results from the GET UP! Study. *Pediatr Obes.* 2019;14(11):e12557.
103. Deng X, He M, He D, Zhu Y, Zhang Z, Niu W. Sleep duration and obesity in children and adolescents: evidence from an updated and dose-response meta-analysis. *Sleep Med.* 2021;78:169-81.
104. Sparano S, Lauria F, Ahrens W, Fraterman A, Thumann B, Iacoviello L, et al. Sleep

- duration and blood pressure in children: Analysis of the pan-European IDEFICS cohort. *J Clin Hypertens (Greenwich)*. 2019;21(5):572-8.
105. Collings PJ, Blackwell JE, Pal E, Ball HL, Wright J. Associations of diarised sleep onset time, period and duration with total and central adiposity in a biethnic sample of young children: the Born in Bradford observational cohort study. *BMJ Open*. 2021;11(5):e044769.
 106. Fiese BH, Cai T, Sutter C, Bost KK. Bedtimes, bedtime routines, and children's sleep across the first 2 years of life. *Sleep*. 2021;44(8).
 107. Chindamo S, Buja A, DeBattisti E, Terraneo A, Marini E, Gomez Perez LJ, et al. Sleep and new media usage in toddlers. *Eur J Pediatr*. 2019;178(4):483-90.
 108. Staples AD, Hoyniak C, McQuillan ME, Molfese V, Bates JE. Screen use before bedtime: Consequences for nighttime sleep in young children. *Infant Behav Dev*. 2021;62:101522.
 109. Bagley EJ, Kelly RJ, Buckhalt JA, El-Sheikh M. What keeps low-SES children from sleeping well: the role of presleep worries and sleep environment. *Sleep Med*. 2015;16(4):496-502.
 110. Hoyniak CP, Bates JE, McQuillan ME, Albert LE, Staples AD, Molfese VJ, et al. The Family Context of Toddler Sleep: Routines, Sleep Environment, and Emotional Security Induction in the Hour before Bedtime. *Behav Sleep Med*. 2021;19(6):795-813.
 111. Perez-Escamilla R, Jimenez EY, Dewey KG. Responsive Feeding Recommendations: Harmonizing Integration into Dietary Guidelines for Infants and Young Children. *Curr Dev Nutr*. 2021;5(6):nzab076.
 112. Pérez-Escamilla R, Segura-Pérez S, Lott M. Feeding guidelines for infants and young toddlers: a responsive parenting approach 2017. 223-31 p.
 113. Vos MB, Kaar JL, Welsh JA, Van Horn LV, Feig DI, Anderson CAM, et al. Added Sugars and Cardiovascular Disease Risk in Children: A Scientific Statement From the American Heart Association. *Circulation*. 2017;135(19):e1017-e34.
 114. Veldman SLC, Chin APMJM, Altenburg TM. Physical activity and prospective associations with indicators of health and development in children aged <5 years: a systematic review. *Int J Behav Nutr Phys Act*. 2021;18(1):6.
 115. Pate RR, Hillman CH, Janz KF, Katzmarzyk PT, Powell KE, Torres A, et al. Physical Activity and Health in Children Younger than 6 Years: A Systematic Review. *Med Sci Sports Exerc*. 2019;51(6):1282-91.
 116. Barnett LM, van Beurden E, Morgan PJ, Brooks LO, Beard JR. Childhood motor skill proficiency as a predictor of adolescent physical activity. *J Adolesc Health*. 2009;44(3):252-9.
 117. Barnett LM, Lai SK, Veldman SLC, Hardy LL, Cliff DP, Morgan PJ, et al. Correlates of Gross Motor Competence in Children and Adolescents: A Systematic Review and Meta-Analysis. *Sports Med*. 2016;46(11):1663-88.
 118. Rudd JR, Crotti M, Fitton-Davies K, O'Callaghan L, Bardid F, Utesch T, et al. Skill Acquisition Methods Fostering Physical Literacy in Early-Physical Education (SAMPLE-

- PE): Rationale and Study Protocol for a Cluster Randomized Controlled Trial in 5-6-Year-Old Children From Deprived Areas of North West England. *Front Psychol.* 2020;11:1228.
119. Mukherjee S, Ting Jamie LC, Fong LH. Fundamental Motor Skill Proficiency of 6- to 9-Year-Old Singaporean Children. *Percept Mot Skills.* 2017;124(3):584-600.
120. Xiong S, Sankaridurg P, Naduvilath T, Zang J, Zou H, Zhu J, et al. Time spent in outdoor activities in relation to myopia prevention and control: a meta-analysis and systematic review. *Acta Ophthalmol.* 2017;95(6):551-66.
121. Lingham G, Yazar S, Lucas RM, Milne E, Hewitt AW, Hammond CJ, et al. Time spent outdoors in childhood is associated with reduced risk of myopia as an adult. *Sci Rep.* 2021;11(1):6337.
122. Early Years Physical Literacy. Physical Literacy Proof of Concept Study in Child Care Settings Canada2020 [Available from: https://afdb32fd-877e-4e25-828f-3b88265b25d2.filesusr.com/uqd/05c80a_2abd1c9b0afd47758303ac3ee9f89008.pdf].
123. Ginsburg KR, American Academy of Pediatrics Committee on C, American Academy of Pediatrics Committee on Psychosocial Aspects of C, Family H. The importance of play in promoting healthy child development and maintaining strong parent-child bonds. *Pediatrics.* 2007;119(1):182-91.
124. Bento G, Dias G. The importance of outdoor play for young children's healthy development. *Porto Biomed J.* 2017;2(5):157-60.
125. Bull FC, Al-Ansari SS, Biddle S, Borodulin K, Buman MP, Cardon G, et al. World Health Organization 2020 guidelines on physical activity and sedentary behaviour. *Br J Sports Med.* 2020;54(24):1451-62.
126. Padmapriya N, Tint MT, Sadananthan SA, Michael N, Chen B, Cai S, et al. The longitudinal association between early-life screen viewing and abdominal adiposity-findings from a multiethnic birth cohort study. *Int J Obes (Lond).* 2021;45(9):1995-2005.
127. Padmapriya N, Aris IM, Tint MT, Loy SL, Cai S, Tan KH, et al. Sex-specific longitudinal associations of screen viewing time in children at 2-3 years with adiposity at 3-5 years. *Int J Obes (Lond).* 2019;43(7):1334-43.
128. McManus AM. Physical activity - a neat solution to an impending crisis. *J Sports Sci Med.* 2007;6(3):368-73.
129. Agras WS, Hammer LD, McNicholas F, Kraemer HC. Risk factors for childhood overweight: a prospective study from birth to 9.5 years. *J Pediatr.* 2004;145(1):20-5.
130. Touchette E, Petit D, Tremblay RE, Boivin M, Falissard B, Genolini C, et al. Associations between sleep duration patterns and overweight/obesity at age 6. *Sleep.* 2008;31(11):1507-14.
131. Diethelm K, Bolzenius K, Cheng G, Remer T, Buyken AE. Longitudinal associations between reported sleep duration in early childhood and the development of body mass index, fat mass index and fat free mass index until age 7. *Int J Pediatr Obes.* 2011;6(2-2):e114-23.
132. Scharf RJ, DeBoer MD. Sleep timing and longitudinal weight gain in 4- and 5-year-old

- children. *Pediatr Obes*. 2015;10(2):141-8.
133. Zhang Z, Adamo KB, Ogden N, Goldfield GS, Okely AD, Kuzik N, et al. Associations between sleep duration, adiposity indicators, and cognitive development in young children. *Sleep Med*. 2021;82:54-60.
 134. Keefe-Cooperman K, Brady-Amoon P. Preschooler sleep patterns related to cognitive and adaptive functioning. *Early Education and Development*. 2014;25(6):859-74.
 135. Vaughn BE, Elmore-Staton L, Shin N, El-Sheikh M. Sleep as a support for social competence, peer relations, and cognitive functioning in preschool children. *Behav Sleep Med*. 2015;13(2):92-106.
 136. Magee C, Caputi P, Iverson D. Lack of sleep could increase obesity in children and too much television could be partly to blame. *Acta Paediatr*. 2014;103(1):e27-31.
 137. Plancoulaine S, Lioret S, Regnault N, Heude B, Charles MA, Eden Mother-Child Cohort Study G. Gender-specific factors associated with shorter sleep duration at age 3 years. *J Sleep Res*. 2015;24(6):610-20.
 138. Ikeda M, Kaneita Y, Kondo S, Itani O, Ohida T. Epidemiological study of sleep habits among four-and-a-half-year-old children in Japan. *Sleep Med*. 2012;13(7):787-94.
 139. Koulouglioti C, Cole R, Kitzman H. Inadequate sleep and unintentional injuries in young children. *Public Health Nurs*. 2008;25(2):106-14.
 140. Boto LR, Crispim JN, de Melo IS, Juvandes C, Rodrigues T, Azeredo P, et al. Sleep deprivation and accidental fall risk in children. *Sleep Med*. 2012;13(1):88-95.
 141. Giganti F, Arzilli C, Conte F, Toselli M, Viggiano MP, Ficca G. The effect of a daytime nap on priming and recognition tasks in preschool children. *Sleep*. 2014;37(6):1087-93.
 142. Bernier A, Beauchamp MH, Bouvette-Turcot AA, Carlson SM, Carrier J. Sleep and cognition in preschool years: specific links to executive functioning. *Child Dev*. 2013;84(5):1542-53.
 143. Jung E, Molfese VJ, Beswick J, Jacobi-Vessels J, Molnar A. Growth of cognitive skills in preschoolers: impact of sleep habits and learning-related behaviors. *Early Education and Development*. 2009;20(4):713-31.
 144. Lam JC, Mahone EM, Mason T, Scharf SM. The effects of napping on cognitive function in preschoolers. *J Dev Behav Pediatr*. 2011;32(2):90-7.
 145. Reynaud E, Forhan A, Heude B, Charles MA, Plancoulaine S. Night-sleep Duration Trajectories and Behavior in Preschoolers: Results from a Prospective Birth Cohort Study. *Behav Sleep Med*. 2021;19(4):445-57.
 146. Dev DA, McBride BA, Fiese BH, Jones BL, Cho H, Behalf Of The Strong Kids Research T. Risk factors for overweight/obesity in preschool children: an ecological approach. *Child Obes*. 2013;9(5):399-408.
 147. Wang H, Sekine M, Chen X, Yamagami T, Kagamimori S. Lifestyle at 3 years of age and quality of life (QOL) in first-year junior high school students in Japan: results of the Toyama Birth Cohort Study. *Qual Life Res*. 2008;17(2):257-65.

148. Scaglioni S, De Cosmi V, Ciappolino V, Parazzini F, Brambilla P, Agostoni C. Factors Influencing Children's Eating Behaviours. *Nutrients*. 2018;10(6).
149. Wellington Ministry of Health. Food and Nutrition Guidelines for Healthy Children and Young People (Aged 2–18 years): A background paper. Partial Revision ed. Wellington: Ministry of Health; 2015 July.
150. Santos R, Zhang Z, Pereira JR, Sousa-Sa E, Cliff DP, Okely AD. Compliance with the Australian 24-hour movement guidelines for the early years: associations with weight status. *BMC Public Health*. 2017;17(Suppl 5):867.
151. Carson V, Tremblay MS, Chastin SFM. Cross-sectional associations between sleep duration, sedentary time, physical activity, and adiposity indicators among Canadian preschool-aged children using compositional analyses. *BMC Public Health*. 2017;17(Suppl 5):848.
152. Kuzik N, Naylor PJ, Spence JC, Carson V. Movement behaviours and physical, cognitive, and social-emotional development in preschool-aged children: Cross-sectional associations using compositional analyses. *PLoS One*. 2020;15(8):e0237945.

Glossary

| | |
|--|--|
| Cardiometabolic health | The interplay of blood pressure, blood lipids, glucose and insulin on health. |
| Cognitive development | The process of learning, memory, attention, concentration and language development. |
| Emotional regulation | An individual's ability to manage and respond to emotional experiences such as stress, anxiety, mood, temperament, hyperactivity/impulsivity. |
| Energetic play | Active play that is equivalent to moderate-to-vigorous physical activity, when children get out of breath and feel warm. This may take many forms and may involve other children, caregivers, objects or not. |
| Exercise | Physical activity that is planned, structured, generally repetitive and has purpose. |
| Light-intensity physical activity | LPA is equivalent to 1.5–4 METs in children, i.e., activities with energy cost 1.5 to 4.0 times the energy expenditure at rest for that child. For young children, this can include slow walking, bathing, or other incidental activities that do not result in the child getting hot or short of breath. |
| Metabolic equivalent of task | The metabolic equivalent of task, or simply metabolic equivalent, is a physiological measure expressing the energy cost (or calories) of physical activities. One MET is the energy equivalent expended by an individual while seated at rest. |

| | |
|--|--|
| <p>Moderate to vigorous intensity physical activity</p> | <p>Moderate PA is equivalent to 4–7 METs in children, i.e., 4–7 times resting energy expenditure at rest for that child.</p> <p>Vigorous PA is equivalent to >7 METs.</p> <p>For young children, this can include brisk walking, cycling, running playing ball games, swimming, dancing etc. during which the child gets hot and breathless.</p> |
| <p>Physical activity</p> | <p>Movement of the body that uses energy over and above resting.</p> <p>For young children, this can include walking, crawling, running, jumping, balancing, climbing in, through and over objects, dancing, riding wheeled toys, cycling, jumping rope etc.</p> |
| <p>Sedentary behaviour</p> | <p>Any waking behaviour characterized by an energy expenditure ≤ 1.5 metabolic equivalents (METs), while in a sitting, reclining or lying posture.</p> <p>For children under 5 years of age includes time spent restrained in car seat, high-chair, stroller, pram or in a carrying device or on a caregiver’s back. Includes time spent sitting quietly listening to a story.</p> |
| <p>Tummy time</p> | <p>Time an infant spends lying on their front (in prone position) while awake with unrestricted movement of limbs.</p> |

Source: Guidelines on Physical Activity, Sedentary Behaviour and Sleep for Children under 5 Years of Age. Geneva: World Health Organization; 2019.

PUBLISHED: January 2022

College of Paediatrics & Child Health, Singapore
Academy of Medicine, Singapore
81 Kim Keat Road
#11-00 NKF Centre
Singapore 328836